



Lecture Tue.2

Many-body methods for electron-phonon interactions

Feliciano Giustino

Oden Institute & Department of Physics
The University of Texas at Austin

Lecture Summary

- Limitations of Rayleigh-Schrödinger perturbation theory
- Many-body Hamiltonian
- Green's function and the spectral function
- Electron-phonon self-energy
- Mass enhancement and electron lifetimes

Limitations of Rayleigh-Schrödinger perturbation theory

Kohn-Sham equations again

$$-\frac{\hbar^2}{2m_e} \nabla^2 \psi_n(\mathbf{r}) + V_{\text{SCF}}(\mathbf{r}; \boldsymbol{\tau}_1, \boldsymbol{\tau}_2, \dots) \psi_n(\mathbf{r}) = \varepsilon_n \psi_n(\mathbf{r})$$

Limitations of Rayleigh-Schrödinger perturbation theory

Kohn-Sham equations again

$$-\frac{\hbar^2}{2m_e} \nabla^2 \psi_n(\mathbf{r}) + V_{\text{SCF}}(\mathbf{r}; \boldsymbol{\tau}_1, \boldsymbol{\tau}_2, \dots) \psi_n(\mathbf{r}) = \varepsilon_n \psi_n(\mathbf{r})$$

- Adiabatic Born-Oppenheimer approximation

Limitations of Rayleigh-Schrödinger perturbation theory

Kohn-Sham equations again

$$-\frac{\hbar^2}{2m_e} \nabla^2 \psi_n(\mathbf{r}) + V_{\text{SCF}}(\mathbf{r}; \boldsymbol{\tau}_1, \boldsymbol{\tau}_2, \dots) \psi_n(\mathbf{r}) = \varepsilon_n \psi_n(\mathbf{r})$$

- Adiabatic Born-Oppenheimer approximation
- Nuclei described as classical particles

Limitations of Rayleigh-Schrödinger perturbation theory

Kohn-Sham equations again

$$-\frac{\hbar^2}{2m_e} \nabla^2 \psi_n(\mathbf{r}) + V_{\text{SCF}}(\mathbf{r}; \boldsymbol{\tau}_1, \boldsymbol{\tau}_2, \dots) \psi_n(\mathbf{r}) = \varepsilon_n \psi_n(\mathbf{r})$$

- Adiabatic Born-Oppenheimer approximation
- Nuclei described as classical particles
- Electron-phonon interactions depend on the XC functional

Breakdown of Rayleigh-Schrödinger perturbation theory

Polaron liquid at the SrTiO₃(001) surface





Figure from Wang et al, Nature Mater. 15, 835 (2016)

Breakdown of Rayleigh-Schrödinger perturbation theory

cf. Lec. Wed.2 Margine

Scanning tunneling spectra of superconducting 2H-NbS₂



Figures from Guillamón et al, Phys. Rev. Lett. 101, 166407 (2008)
and Heil et al, Phys. Rev. Lett. 119, 087003 (2017)

Many-body Schrödinger equation for electrons and nuclei

$$\left[-\frac{\hbar^2}{2m_e} \sum_i \nabla_i^2 - \frac{\hbar^2}{2M_\kappa} \sum_\kappa \nabla_\kappa^2 - \sum_{i,\kappa} Z_\kappa v(\mathbf{r}_i, \boldsymbol{\tau}_\kappa) + \sum_{\kappa > \kappa'} Z_\kappa Z_{\kappa'} v(\boldsymbol{\tau}_\kappa, \boldsymbol{\tau}_{\kappa'}) + \sum_{i>j} v(\mathbf{r}_i, \mathbf{r}_j) \right] \Psi = E_{\text{tot}} \Psi$$

$$\mathbf{r}_i \text{ electron}, \boldsymbol{\tau}_\kappa \text{ nucleus}, v(\mathbf{r}, \mathbf{r}') = \frac{e^2}{4\pi\epsilon_0 |\mathbf{r} - \mathbf{r}'|}$$

Many-body Schrödinger equation for electrons and nuclei

$$\left[-\frac{\hbar^2}{2m_e} \sum_i \nabla_i^2 - \frac{\hbar^2}{2M_\kappa} \sum_\kappa \nabla_\kappa^2 - \sum_{i,\kappa} Z_\kappa v(\mathbf{r}_i, \boldsymbol{\tau}_\kappa) + \sum_{\kappa > \kappa'} Z_\kappa Z_{\kappa'} v(\boldsymbol{\tau}_\kappa, \boldsymbol{\tau}_{\kappa'}) + \sum_{i>j} v(\mathbf{r}_i, \mathbf{r}_j) \right] \Psi = E_{\text{tot}} \Psi$$

$$\mathbf{r}_i \text{ electron}, \boldsymbol{\tau}_\kappa \text{ nucleus}, v(\mathbf{r}, \mathbf{r}') = \frac{e^2}{4\pi\epsilon_0 |\mathbf{r} - \mathbf{r}'|}$$

- Electrons and vibrations must be described on the same footing

Many-body Schrödinger equation for electrons and nuclei

$$\left[-\frac{\hbar^2}{2m_e} \sum_i \nabla_i^2 - \frac{\hbar^2}{2M_\kappa} \sum_\kappa \nabla_\kappa^2 - \sum_{i,\kappa} Z_\kappa v(\mathbf{r}_i, \boldsymbol{\tau}_\kappa) + \sum_{\kappa > \kappa'} Z_\kappa Z_{\kappa'} v(\boldsymbol{\tau}_\kappa, \boldsymbol{\tau}_{\kappa'}) + \sum_{i>j} v(\mathbf{r}_i, \mathbf{r}_j) \right] \Psi = E_{\text{tot}} \Psi$$

$$\mathbf{r}_i \text{ electron}, \boldsymbol{\tau}_\kappa \text{ nucleus}, v(\mathbf{r}, \mathbf{r}') = \frac{e^2}{4\pi\epsilon_0 |\mathbf{r} - \mathbf{r}'|}$$

- Electrons and vibrations must be described on the same footing
- The many-body Schrödinger equation is impractical for calculations or EPIs

Field operators

Many-electron wavefunction as a linear combination of Slater determinants

$$\Psi(\mathbf{r}_1, \mathbf{r}_2, \dots) = \sum_{mn} A_{mn} \hat{c}_m^\dagger \hat{c}_n |0_{\text{KS}}\rangle + \sum_{mnpq} B_{mnpq} \hat{c}_m^\dagger \hat{c}_n^\dagger \hat{c}_p \hat{c}_q |0_{\text{KS}}\rangle + \dots$$

Field operators

Many-electron wavefunction as a linear combination of Slater determinants

$$\Psi(\mathbf{r}_1, \mathbf{r}_2, \dots) = \sum_{mn} A_{mn} \hat{c}_m^\dagger \hat{c}_n |0_{\text{KS}}\rangle + \sum_{mnpq} B_{mnpq} \hat{c}_m^\dagger \hat{c}_n^\dagger \hat{c}_p \hat{c}_q |0_{\text{KS}}\rangle + \dots$$

Operators in second quantization

$$\sum_i V(\mathbf{r}_i) \longrightarrow \sum_{mn} V_{mn} \hat{c}_m^\dagger \hat{c}_n$$

Field operators

Many-electron wavefunction as a linear combination of Slater determinants

$$\Psi(\mathbf{r}_1, \mathbf{r}_2, \dots) = \sum_{mn} A_{mn} \hat{c}_m^\dagger \hat{c}_n |0_{\text{KS}}\rangle + \sum_{mnpq} B_{mnpq} \hat{c}_m^\dagger \hat{c}_n^\dagger \hat{c}_p \hat{c}_q |0_{\text{KS}}\rangle + \dots$$

Operators in second quantization

$$\sum_i V(\mathbf{r}_i) \longrightarrow \sum_{mn} V_{mn} \hat{c}_m^\dagger \hat{c}_n$$

$$\sum_i V(\mathbf{r}_i) = \sum_m \sum_n \int d\mathbf{r} \psi_m^*(\mathbf{r}) V(\mathbf{r}) \psi_n(\mathbf{r}) \hat{c}_m^\dagger \hat{c}_n$$

Field operators

Many-electron wavefunction as a linear combination of Slater determinants

$$\Psi(\mathbf{r}_1, \mathbf{r}_2, \dots) = \sum_{mn} A_{mn} \hat{c}_m^\dagger \hat{c}_n |0_{\text{KS}}\rangle + \sum_{mnpq} B_{mnpq} \hat{c}_m^\dagger \hat{c}_n^\dagger \hat{c}_p \hat{c}_q |0_{\text{KS}}\rangle + \dots$$

Operators in second quantization

$$\sum_i V(\mathbf{r}_i) \longrightarrow \sum_{mn} V_{mn} \hat{c}_m^\dagger \hat{c}_n$$

$$\sum_i V(\mathbf{r}_i) = \sum_m \sum_n \int d\mathbf{r} \psi_m^*(\mathbf{r}) V(\mathbf{r}) \psi_n(\mathbf{r}) \hat{c}_m^\dagger \hat{c}_n$$

Field operators

$$\hat{\psi}^\dagger(\mathbf{r}) \stackrel{\text{def}}{=} \sum_m \psi_m^*(\mathbf{r}) \hat{c}_m^\dagger \quad \hat{\psi}(\mathbf{r}) \stackrel{\text{def}}{=} \sum_n \psi_n(\mathbf{r}) \hat{c}_n$$

Field operators

Many-electron wavefunction as a linear combination of Slater determinants

$$\Psi(\mathbf{r}_1, \mathbf{r}_2, \dots) = \sum_{mn} A_{mn} \hat{c}_m^\dagger \hat{c}_n |0_{\text{KS}}\rangle + \sum_{mnpq} B_{mnpq} \hat{c}_m^\dagger \hat{c}_n^\dagger \hat{c}_p \hat{c}_q |0_{\text{KS}}\rangle + \dots$$

Operators in second quantization

$$\sum_i V(\mathbf{r}_i) \longrightarrow \sum_{mn} V_{mn} \hat{c}_m^\dagger \hat{c}_n$$

$$\sum_i V(\mathbf{r}_i) = \sum_m \sum_n \int d\mathbf{r} \psi_m^*(\mathbf{r}) V(\mathbf{r}) \psi_n(\mathbf{r}) \hat{c}_m^\dagger \hat{c}_n = \int d\mathbf{r} \hat{\psi}^\dagger(\mathbf{r}) V(\mathbf{r}) \hat{\psi}(\mathbf{r})$$

Field operators

$$\hat{\psi}^\dagger(\mathbf{r}) \stackrel{\text{def}}{=} \sum_m \psi_m^*(\mathbf{r}) \hat{c}_m^\dagger \quad \hat{\psi}(\mathbf{r}) \stackrel{\text{def}}{=} \sum_n \psi_n(\mathbf{r}) \hat{c}_n$$

Hamiltonian in field-theoretic formulation

$$\hat{H} = \hat{T}_e + \hat{T}_n + \hat{U}_{en} + \hat{U}_{ee} + \hat{U}_{nn}$$

$$\hat{n}_e(\mathbf{r}) = \hat{\psi}^\dagger(\mathbf{r})\hat{\psi}(\mathbf{r})$$

Hamiltonian in field-theoretic formulation

$$\hat{H} = \hat{T}_e + \hat{T}_n + \hat{U}_{en} + \hat{U}_{ee} + \hat{U}_{nn}$$

$$\hat{n}_e(\mathbf{r}) = \hat{\psi}^\dagger(\mathbf{r})\hat{\psi}(\mathbf{r})$$

Electron-nucleus interaction $\hat{U}_{en} = \int d\mathbf{r} d\mathbf{r}' \hat{n}_e(\mathbf{r}) \hat{n}_n(\mathbf{r}') v(\mathbf{r}, \mathbf{r}')$

Hamiltonian in field-theoretic formulation

$$\hat{H} = \hat{T}_e + \hat{T}_n + \hat{U}_{en} + \hat{U}_{ee} + \hat{U}_{nn}$$

$$\hat{n}_e(\mathbf{r}) = \hat{\psi}^\dagger(\mathbf{r})\hat{\psi}(\mathbf{r})$$

Electron-nucleus interaction $\hat{U}_{en} = \int d\mathbf{r} d\mathbf{r}' \hat{n}_e(\mathbf{r}) \hat{n}_n(\mathbf{r}') v(\mathbf{r}, \mathbf{r}')$

Electron-electron interaction $\hat{U}_{ee} = \frac{1}{2} \int d\mathbf{r} d\mathbf{r}' \hat{n}_e(\mathbf{r}) [\hat{n}_e(\mathbf{r}') - \delta(\mathbf{r} - \mathbf{r}')] v(\mathbf{r}, \mathbf{r}')$

Time evolution of field operators and Dyson orbitals

Ground state of N -electron system

$$\hat{H}|N\rangle = E_N|N\rangle$$

Time evolution of field operators and Dyson orbitals

Ground state of N -electron system

$$\hat{H}|N\rangle = E_N|N\rangle$$

s -th excited state of $N+1$ -electron system

$$\hat{H}|N + 1, s\rangle = E_{N+1,s}|N + 1, s\rangle$$

Time evolution of field operators and Dyson orbitals

Ground state of N -electron system

$$\hat{H}|N\rangle = E_N|N\rangle$$

s -th excited state of $N+1$ -electron system

$$\hat{H}|N + 1, s\rangle = E_{N+1,s}|N + 1, s\rangle$$

Excitation energy

$$\varepsilon_s = E_{N+1,s} - E_N$$

Time evolution of field operators and Dyson orbitals

Ground state of N -electron system

$$\hat{H}|N\rangle = E_N|N\rangle$$

s -th excited state of $N+1$ -electron system

$$\hat{H}|N + 1, s\rangle = E_{N+1,s}|N + 1, s\rangle$$

Excitation energy

$$\varepsilon_s = E_{N+1,s} - E_N$$

Heisenberg time evolution

$$\hat{\psi}(\mathbf{x}, t) = e^{i\hat{H}t/\hbar} \hat{\psi}(\mathbf{x}) e^{-i\hat{H}t/\hbar}$$

Time evolution of field operators and Dyson orbitals

Ground state of N -electron system

$$\hat{H}|N\rangle = E_N|N\rangle$$

s -th excited state of $N+1$ -electron system

$$\hat{H}|N + 1, s\rangle = E_{N+1,s}|N + 1, s\rangle$$

Excitation energy

$$\varepsilon_s = E_{N+1,s} - E_N$$

Heisenberg time evolution

$$\hat{\psi}(\mathbf{x}, t) = e^{i\hat{H}t/\hbar} \hat{\psi}(\mathbf{x}) e^{-i\hat{H}t/\hbar}$$

Exercise

$$\langle N | \hat{\psi}(\mathbf{x}, t) | N + 1, s \rangle$$

Time evolution of field operators and Dyson orbitals

Ground state of N -electron system

$$\hat{H}|N\rangle = E_N|N\rangle$$

s -th excited state of $N+1$ -electron system

$$\hat{H}|N + 1, s\rangle = E_{N+1,s}|N + 1, s\rangle$$

Excitation energy

$$\varepsilon_s = E_{N+1,s} - E_N$$

Heisenberg time evolution

$$\hat{\psi}(\mathbf{x}, t) = e^{i\hat{H}t/\hbar} \hat{\psi}(\mathbf{x}) e^{-i\hat{H}t/\hbar}$$

Exercise

$$\langle N|\hat{\psi}(\mathbf{x}, t)|N + 1, s\rangle = \langle N|e^{i\hat{H}t/\hbar} \hat{\psi}(\mathbf{x}) e^{-i\hat{H}t/\hbar}|N + 1, s\rangle$$

Time evolution of field operators and Dyson orbitals

Ground state of N -electron system

$$\hat{H}|N\rangle = E_N|N\rangle$$

s -th excited state of $N+1$ -electron system

$$\hat{H}|N + 1, s\rangle = E_{N+1,s}|N + 1, s\rangle$$

Excitation energy

$$\varepsilon_s = E_{N+1,s} - E_N$$

Heisenberg time evolution

$$\hat{\psi}(\mathbf{x}, t) = e^{i\hat{H}t/\hbar} \hat{\psi}(\mathbf{x}) e^{-i\hat{H}t/\hbar}$$

Exercise

$$\begin{aligned}\langle N | \hat{\psi}(\mathbf{x}, t) | N + 1, s \rangle &= \langle N | e^{i\hat{H}t/\hbar} \hat{\psi}(\mathbf{x}) e^{-i\hat{H}t/\hbar} | N + 1, s \rangle \\ &= \langle N | e^{iE_N t/\hbar} \hat{\psi}(\mathbf{x}) e^{-iE_{N+1,s} t/\hbar} | N + 1, s \rangle\end{aligned}$$

Time evolution of field operators and Dyson orbitals

Ground state of N -electron system

$$\hat{H}|N\rangle = E_N|N\rangle$$

s -th excited state of $N+1$ -electron system

$$\hat{H}|N+1, s\rangle = E_{N+1,s}|N+1, s\rangle$$

Excitation energy

$$\varepsilon_s = E_{N+1,s} - E_N$$

Heisenberg time evolution

$$\hat{\psi}(\mathbf{x}, t) = e^{i\hat{H}t/\hbar} \hat{\psi}(\mathbf{x}) e^{-i\hat{H}t/\hbar}$$

Exercise

$$\begin{aligned}\langle N | \hat{\psi}(\mathbf{x}, t) | N + 1, s \rangle &= \langle N | e^{i\hat{H}t/\hbar} \hat{\psi}(\mathbf{x}) e^{-i\hat{H}t/\hbar} | N + 1, s \rangle \\ &= \langle N | e^{iE_N t/\hbar} \hat{\psi}(\mathbf{x}) e^{-iE_{N+1,s} t/\hbar} | N + 1, s \rangle \\ &= \langle N | \hat{\psi}(\mathbf{x}) | N + 1, s \rangle e^{-i\varepsilon_s t/\hbar}\end{aligned}$$

Time evolution of field operators and Dyson orbitals

Ground state of N -electron system

$$\hat{H}|N\rangle = E_N|N\rangle$$

s -th excited state of $N+1$ -electron system

$$\hat{H}|N+1, s\rangle = E_{N+1,s}|N+1, s\rangle$$

Excitation energy

$$\varepsilon_s = E_{N+1,s} - E_N$$

Heisenberg time evolution

$$\hat{\psi}(\mathbf{x}, t) = e^{i\hat{H}t/\hbar} \hat{\psi}(\mathbf{x}) e^{-i\hat{H}t/\hbar}$$

Exercise

$$\begin{aligned}\langle N | \hat{\psi}(\mathbf{x}, t) | N + 1, s \rangle &= \langle N | e^{i\hat{H}t/\hbar} \hat{\psi}(\mathbf{x}) e^{-i\hat{H}t/\hbar} | N + 1, s \rangle \\ &= \langle N | e^{iE_N t/\hbar} \hat{\psi}(\mathbf{x}) e^{-iE_{N+1,s} t/\hbar} | N + 1, s \rangle \\ &= \underbrace{\langle N | \hat{\psi}(\mathbf{x}) | N + 1, s \rangle}_{f_s(\mathbf{x})} e^{-i\varepsilon_s t/\hbar}\end{aligned}$$

Dyson orbital

The Green's function at zero temperature

Time-ordered
Green's function

Wick's time-ordering operator

$$G(\mathbf{x}t, \mathbf{x}'t') = -\frac{i}{\hbar} \langle N | \hat{T} \hat{\psi}(\mathbf{x}t) \hat{\psi}^\dagger(\mathbf{x}'t') | N \rangle$$

The Green's function at zero temperature

Time-ordered
Green's function

$$G(\mathbf{x}t, \mathbf{x}'t') = -\frac{i}{\hbar} \langle N | \hat{T} \hat{\psi}(\mathbf{x}t) \hat{\psi}^\dagger(\mathbf{x}'t') | N \rangle$$

Wick's time-ordering operator

electron in \mathbf{x}' at time t'

The Green's function at zero temperature

Time-ordered
Green's function

$$G(\mathbf{x}t, \mathbf{x}'t') = -\frac{i}{\hbar} \langle N | \hat{T} \hat{\psi}(\mathbf{x}t) \hat{\psi}^\dagger(\mathbf{x}'t') | N \rangle$$

Wick's time-ordering operator

$\left\langle \begin{array}{c|c} \text{electron in } \mathbf{x} \text{ at time } t & \text{electron in } \mathbf{x}' \text{ at time } t' \end{array} \right\rangle$

The Green's function at zero temperature

Time-ordered
Green's function

$$G(\mathbf{x}t, \mathbf{x}'t') = -\frac{i}{\hbar} \langle N | \hat{T} \hat{\psi}(\mathbf{x}t) \hat{\psi}^\dagger(\mathbf{x}'t') | N \rangle$$

Wick's time-ordering operator

\langle electron in \mathbf{x} at time t | electron in \mathbf{x}' at time t' \rangle



The Green's function at zero temperature

Consider $t > t'$ (electron added to ground state)

$$G(\mathbf{x}t, \mathbf{x}'t') = -\frac{i}{\hbar} \langle N | \hat{\psi}(\mathbf{x}t) \hat{\psi}^\dagger(\mathbf{x}'t') | N \rangle$$

The Green's function at zero temperature

Consider $t > t'$ (electron added to ground state)

$$\begin{aligned} G(\mathbf{x}t, \mathbf{x}'t') &= -\frac{i}{\hbar} \langle N | \hat{\psi}(\mathbf{x}t) \hat{\psi}^\dagger(\mathbf{x}'t') | N \rangle \\ &= -\frac{i}{\hbar} \langle N | e^{i\hat{H}t/\hbar} \hat{\psi}(\mathbf{x}) e^{-i\hat{H}t/\hbar} e^{i\hat{H}t'/\hbar} \hat{\psi}^\dagger(\mathbf{x}') e^{-i\hat{H}t'/\hbar} | N \rangle \end{aligned}$$

The Green's function at zero temperature

Consider $t > t'$ (electron added to ground state)

$$\begin{aligned} G(\mathbf{x}t, \mathbf{x}'t') &= -\frac{i}{\hbar} \langle N | \hat{\psi}(\mathbf{x}t) \hat{\psi}^\dagger(\mathbf{x}'t') | N \rangle \\ &= -\frac{i}{\hbar} \langle N | e^{i\hat{H}t/\hbar} \hat{\psi}(\mathbf{x}) e^{-i\hat{H}t/\hbar} e^{i\hat{H}t'/\hbar} \hat{\psi}^\dagger(\mathbf{x}') e^{-i\hat{H}t'/\hbar} | N \rangle \\ &= -\frac{i}{\hbar} \langle N | \hat{\psi}(\mathbf{x}) e^{-i(\hat{H}-E_N)(t-t')/\hbar} \hat{\psi}^\dagger(\mathbf{x}') | N \rangle \end{aligned}$$

The Green's function at zero temperature

Consider $t > t'$ (electron added to ground state)

$$\begin{aligned} G(\mathbf{x}t, \mathbf{x}'t') &= -\frac{i}{\hbar} \langle N | \hat{\psi}(\mathbf{x}t) \hat{\psi}^\dagger(\mathbf{x}'t') | N \rangle \\ &= -\frac{i}{\hbar} \langle N | e^{i\hat{H}t/\hbar} \hat{\psi}(\mathbf{x}) e^{-i\hat{H}t/\hbar} e^{i\hat{H}t'/\hbar} \hat{\psi}^\dagger(\mathbf{x}') e^{-i\hat{H}t'/\hbar} | N \rangle \\ &= -\frac{i}{\hbar} \langle N | \hat{\psi}(\mathbf{x}) \Big| e^{-i(\hat{H}-E_N)(t-t')/\hbar} \Big| \hat{\psi}^\dagger(\mathbf{x}') | N \rangle \end{aligned}$$

$\sum_s |N+1, s\rangle \langle N+1, s|$

The Green's function at zero temperature

Consider $t > t'$ (electron added to ground state)

$$\begin{aligned} G(\mathbf{x}t, \mathbf{x}'t') &= -\frac{i}{\hbar} \langle N | \hat{\psi}(\mathbf{x}t) \hat{\psi}^\dagger(\mathbf{x}'t') | N \rangle \\ &= -\frac{i}{\hbar} \langle N | e^{i\hat{H}t/\hbar} \hat{\psi}(\mathbf{x}) e^{-i\hat{H}t/\hbar} e^{i\hat{H}t'/\hbar} \hat{\psi}^\dagger(\mathbf{x}') e^{-i\hat{H}t'/\hbar} | N \rangle \\ &= -\frac{i}{\hbar} \langle N | \hat{\psi}(\mathbf{x}) \Big| e^{-i(\hat{H}-E_N)(t-t')/\hbar} \Big| \hat{\psi}^\dagger(\mathbf{x}') | N \rangle \\ &\quad \boxed{\sum_s |N+1, s\rangle \langle N+1, s|} \\ &= -\frac{i}{\hbar} \sum_s f_s(\mathbf{x}) f_s^*(\mathbf{x}') e^{-i\varepsilon_s(t-t')/\hbar} \end{aligned}$$

The spectral function

Carry out the same operation for $t < t'$ and Fourier transform

$$G(\mathbf{x}, \mathbf{x}', \omega) = \sum_s \frac{f_s(\mathbf{x}) f_s^*(\mathbf{x}')}{\hbar\omega - \varepsilon_s \mp i0^+} \quad \mp \text{occ/unocc}$$

The spectral function

Carry out the same operation for $t < t'$ and Fourier transform

$$G(\mathbf{x}, \mathbf{x}', \omega) = \sum_s \frac{f_s(\mathbf{x}) f_s^*(\mathbf{x}')}{\hbar\omega - \varepsilon_s \mp i0^+} \quad \mp \text{occ/unocc}$$

The poles of the Green's function represent the electron addition/removal energies of the interacting many-body system

The spectral function

Carry out the same operation for $t < t'$ and Fourier transform

$$G(\mathbf{x}, \mathbf{x}', \omega) = \sum_s \frac{f_s(\mathbf{x}) f_s^*(\mathbf{x}')}{\hbar\omega - \varepsilon_s \mp i0^+} \quad \mp \text{occ/unocc}$$

The poles of the Green's function represent the electron addition/removal energies of the interacting many-body system

From the Green's function we can obtain the **spectral (density)** function

$$A(\mathbf{x}, \omega) = \frac{1}{\pi} |\text{Im } G(\mathbf{x}, \mathbf{x}, \omega)| = \sum_s |f_s(\mathbf{x})|^2 \delta(\hbar\omega - \varepsilon_s)$$

The spectral function

Carry out the same operation for $t < t'$ and Fourier transform

$$G(\mathbf{x}, \mathbf{x}', \omega) = \sum_s \frac{f_s(\mathbf{x}) f_s^*(\mathbf{x}')}{\hbar\omega - \varepsilon_s \mp i0^+} \quad \mp \text{occ/unocc}$$

The poles of the Green's function represent the electron addition/removal energies of the interacting many-body system

From the Green's function we can obtain the **spectral (density)** function


$$A(\mathbf{x}, \omega) = \frac{1}{\pi} |\text{Im } G(\mathbf{x}, \mathbf{x}, \omega)| = \sum_s |f_s(\mathbf{x})|^2 \delta(\hbar\omega - \varepsilon_s)$$

The spectral function is the many-body (local) density of states

Usually presented as momentum-resolved $A(\mathbf{k}, \omega)$


The spectral function: Broadening

Example: a single complex pole $\varepsilon_s = \varepsilon - i\Gamma$



The spectral function: Broadening


Example: a single complex pole $\varepsilon_s = \varepsilon - i\Gamma$



$$G(\mathbf{x}, \mathbf{x}, t-t') = -\frac{i}{\hbar} |f_s(\mathbf{x})|^2 e^{-i\varepsilon(t-t')/\hbar} \underline{e^{-\Gamma(t-t')/\hbar}}$$

The spectral function: Broadening

Example: a single complex pole $\varepsilon_s = \varepsilon - i\Gamma$



$$G(\mathbf{x}, \mathbf{x}, t-t') = -\frac{i}{\hbar} |f_s(\mathbf{x})|^2 e^{-i\varepsilon(t-t')/\hbar} \underline{e^{-\Gamma(t-t')/\hbar}}$$



$$A(\mathbf{x}, \mathbf{x}, \omega) = \frac{1}{\pi} \frac{\Gamma}{(\hbar\omega - \varepsilon)^2 + \Gamma^2} |f_s(\mathbf{x})|^2$$

The spectral function: Broadening

Example: a single complex pole $\varepsilon_s = \varepsilon - i\Gamma$

$$G(\mathbf{x}, \mathbf{x}, t-t') = -\frac{i}{\hbar} |f_s(\mathbf{x})|^2 e^{-i\varepsilon(t-t')/\hbar} \underline{e^{-\Gamma(t-t')/\hbar}}$$

$$A(\mathbf{x}, \mathbf{x}, \omega) = \frac{1}{\pi} \frac{\Gamma}{(\hbar\omega - \varepsilon)^2 + \Gamma^2} |f_s(\mathbf{x})|^2$$




The spectral function: Coherent and incoherent structures

$$A(\mathbf{k}, \omega) = \frac{1}{\pi} |\text{Im } G(\mathbf{k}, \omega)|$$


The spectral function: Coherent and incoherent structures

$$A(\mathbf{k}, \omega) = \frac{1}{\pi} |\text{Im } G(\mathbf{k}, \omega)|$$




The spectral function: Coherent and incoherent structures

$$A(\mathbf{k}, \omega) = \frac{1}{\pi} |\text{Im } G(\mathbf{k}, \omega)|$$




The spectral function: Coherent and incoherent structures

$$A(\mathbf{k}, \omega) = \frac{1}{\pi} |\text{Im } G(\mathbf{k}, \omega)|$$




The spectral function: Coherent and incoherent structures

$$A(\mathbf{k}, \omega) = \frac{1}{\pi} |\text{Im } G(\mathbf{k}, \omega)|$$



The spectral function: Coherent and incoherent structures

$$A(\mathbf{k}, \omega) = \frac{1}{\pi} |\text{Im } G(\mathbf{k}, \omega)|$$



How to calculate the Green's function

Heisenberg time evolution

$$\hat{\psi}(\mathbf{x}, t) = e^{i\hat{H}t/\hbar} \hat{\psi}(\mathbf{x}) e^{-i\hat{H}t/\hbar}$$

How to calculate the Green's function

Heisenberg time evolution

$$\hat{\psi}(\mathbf{x}, t) = e^{i\hat{H}t/\hbar} \hat{\psi}(\mathbf{x}) e^{-i\hat{H}t/\hbar}$$

Equation of motion for field operators

$$i\hbar \frac{\partial}{\partial t} \hat{\psi}(\mathbf{x}t) = [\hat{\psi}(\mathbf{x}, t), \hat{H}]$$

How to calculate the Green's function

Heisenberg time evolution

$$\hat{\psi}(\mathbf{x}, t) = e^{i\hat{H}t/\hbar} \hat{\psi}(\mathbf{x}) e^{-i\hat{H}t/\hbar}$$

Equation of motion for field operators

$$i\hbar \frac{\partial}{\partial t} \hat{\psi}(\mathbf{x}t) = [\hat{\psi}(\mathbf{x}, t), \hat{H}] = \left[-\frac{\hbar^2}{2m_e} \nabla^2 + \int d\mathbf{r}' v(\mathbf{r}, \mathbf{r}') \hat{n}(\mathbf{r}'t) \right] \hat{\psi}(\mathbf{x}t)$$

total charge, electrons & nuclei 

How to calculate the Green's function

Heisenberg time evolution

$$\hat{\psi}(\mathbf{x}, t) = e^{i\hat{H}t/\hbar} \hat{\psi}(\mathbf{x}) e^{-i\hat{H}t/\hbar}$$

Equation of motion for field operators

$$i\hbar \frac{\partial}{\partial t} \hat{\psi}(\mathbf{x}t) = [\hat{\psi}(\mathbf{x}, t), \hat{H}] = \left[-\frac{\hbar^2}{2m_e} \nabla^2 + \int d\mathbf{r}' v(\mathbf{r}, \mathbf{r}') \hat{n}(\mathbf{r}'t) \right] \hat{\psi}(\mathbf{x}t)$$

total charge, electrons & nuclei 

Equation of motion for Green's function

$$\left[i\hbar \frac{\partial}{\partial t_1} + \frac{\hbar^2}{2m_e} \nabla_1^2 \right] G(12) + \frac{i}{\hbar} \int d3 v(13) \langle \hat{T} \hat{n}(3) \psi(1) \psi^\dagger(2) \rangle = \delta(12)$$

How to calculate the Green's function

Heisenberg time evolution

$$\hat{\psi}(\mathbf{x}, t) = e^{i\hat{H}t/\hbar} \hat{\psi}(\mathbf{x}) e^{-i\hat{H}t/\hbar}$$

Equation of motion for field operators

$$i\hbar \frac{\partial}{\partial t} \hat{\psi}(\mathbf{x}t) = [\hat{\psi}(\mathbf{x}, t), \hat{H}] = \left[-\frac{\hbar^2}{2m_e} \nabla^2 + \int d\mathbf{r}' v(\mathbf{r}, \mathbf{r}') \hat{n}(\mathbf{r}'t) \right] \hat{\psi}(\mathbf{x}t)$$

total charge, electrons & nuclei

Equation of motion for Green's function

$$\left[i\hbar \frac{\partial}{\partial t_1} + \frac{\hbar^2}{2m_e} \nabla_1^2 \right] G(12) + \frac{i}{\hbar} \int d3 v(13) \langle \hat{T} \hat{n}(3) \psi(1) \psi^\dagger(2) \rangle = \delta(12)$$

4 field operators

Hartree+Fock+2-particle Green's function

Dyson equation

$$V_{\text{tot}}(1) = \int d2 v(12) \langle \hat{n}(2) \rangle$$

2-particle Green's function
rewritten using self-energy Σ

$$\left[i\hbar \frac{\partial}{\partial t_1} + \frac{\hbar^2}{2m_e} \nabla_1^2 - V_{\text{tot}}(1) \right] G(12) - \int d3 \Sigma(13) G(32) = \delta(12)$$

Dyson equation

$$V_{\text{tot}}(1) = \int d2 v(12) \langle \hat{n}(2) \rangle \quad \begin{array}{c} \longrightarrow \\ \downarrow \end{array} \quad \begin{array}{l} \text{2-particle Green's function} \\ \text{rewritten using self-energy } \Sigma \end{array}$$
$$\left[i\hbar \frac{\partial}{\partial t_1} + \frac{\hbar^2}{2m_e} \nabla_1^2 - V_{\text{tot}}(1) \right] G(12) - \int d3 \Sigma(13) G(32) = \delta(12)$$

Express the Green's function in terms of Dyson's orbitals

$$\left[-\frac{\hbar^2}{2m_e} \nabla^2 + V_{\text{tot}}(\mathbf{r}) \right] f_s(\mathbf{x}) + \int d\mathbf{x}' \Sigma(\mathbf{x}, \mathbf{x}', \varepsilon_s/\hbar) f_s(\mathbf{x}') = \varepsilon_s f_s(\mathbf{x})$$

Dyson equation

$$V_{\text{tot}}(1) = \int d2 v(12) \langle \hat{n}(2) \rangle \quad \begin{array}{l} \text{2-particle Green's function} \\ \text{rewritten using self-energy } \Sigma \end{array}$$
$$\left[i\hbar \frac{\partial}{\partial t_1} + \frac{\hbar^2}{2m_e} \nabla_1^2 - V_{\text{tot}}(1) \right] G(12) - \int d3 \Sigma(13) G(32) = \delta(12)$$

Express the Green's function in terms of Dyson's orbitals

$$\left[-\frac{\hbar^2}{2m_e} \nabla^2 + \underline{V_{\text{tot}}(\mathbf{r})} \right] f_s(\mathbf{x}) + \int d\mathbf{x}' \underline{\Sigma(\mathbf{x}, \mathbf{x}', \varepsilon_s/\hbar)} f_s(\mathbf{x}') = \varepsilon_s f_s(\mathbf{x})$$

Sources of **electron-phonon** interaction

How to calculate the electron-phonon self-energy

Electron self-energy from Hedin-Baym's equations

$$\Sigma(12) = i\hbar \int d(34) G(13) \Gamma(324) W(41^+)$$

↑
Green's function
↑
Vertex
↑
Screened Coulomb interaction

How to calculate the electron-phonon self-energy

Electron self-energy from Hedin-Baym's equations

$$\Sigma(12) = i\hbar \int d(34) G(13) \Gamma(324) W(41^+)$$

↑
Green's function
↑
Vertex
↑
Screened Coulomb interaction

$$W = W_e + W_{ph}$$

$$W_e(12) = \int d3 \epsilon_e^{-1}(13) v(32)$$

Reduces to the standard GW method + screening from nuclei

Diagrammatic representation of the self-energy




Figure from

FG, Rev. Mod. Phys. 89, 015003 (2017)

Diagrammatic representation of the self-energy




Figure from

FG, Rev. Mod. Phys. 89, 015003 (2017)

Diagrammatic representation of the self-energy




Figure from
FG, Rev. Mod. Phys. 89, 015003 (2017)

Improper self-energy: comes from
 $V_{\text{tot}}(1) = \int d^2 r(1) \langle \hat{n}(2) \rangle$ term

Diagrammatic representation of the self-energy





Figure from
FG, Rev. Mod. Phys. 89, 015003 (2017)

Improper self-energy: comes from
 $V_{\text{tot}}(1) = \int d^2 r(1) \langle \hat{n}(1) \rangle$ term



Fan-Migdal self-energy using Kohn-Sham states and DFPT phonons

$$\Sigma_{n\mathbf{k}}^{\text{FM}}(\omega) = \frac{1}{\hbar} \sum_{m\nu} \int \frac{d\mathbf{q}}{\Omega_{\text{BZ}}} |g_{mn\nu}(\mathbf{k}, \mathbf{q})|^2 \\ \times \left[\frac{1 - f_{m\mathbf{k}+\mathbf{q}} + n_{\mathbf{q}\nu}}{\omega - \varepsilon_{m\mathbf{k}+\mathbf{q}}/\hbar - \omega_{\mathbf{q}\nu} + i\eta} + \frac{f_{m\mathbf{k}+\mathbf{q}} + n_{\mathbf{q}\nu}}{\omega - \varepsilon_{m\mathbf{k}+\mathbf{q}}/\hbar + \omega_{\mathbf{q}\nu} + i\eta} \right]$$



Example: Interaction with dispersionless phonon





Example: Interaction with dispersionless phonon





Example: Interaction with dispersionless phonon





Example: Interaction with dispersionless phonon



Example: Interaction with dispersionless phonon



Example: Interaction with dispersionless phonon



Example from experiments: Velocity renormalization in MgB₂





Figure from Mou et al, Phys. Rev. B 91, 140502(R) (2015)

Quasiparticle shift and broadening

Spectral function from the self-energy

$$A(\mathbf{k}, \omega) = -\frac{1}{\pi} \operatorname{Im} \sum_n \frac{1}{\hbar\omega - \varepsilon_{n\mathbf{k}} - \Sigma_{n\mathbf{k}}(\omega)}$$

Quasiparticle shift and broadening

Spectral function from the self-energy

$$A(\mathbf{k}, \omega) = -\frac{1}{\pi} \operatorname{Im} \sum_n \frac{1}{\hbar\omega - \varepsilon_{n\mathbf{k}} - \Sigma_{n\mathbf{k}}(\omega)}$$

Quasiparticle approximation: assume simple poles in the complex plane

$$\Sigma_{n\mathbf{k}}(\omega) = \Sigma_{n\mathbf{k}}(z) + \frac{1}{\hbar} \left. \frac{\partial \operatorname{Re} \Sigma_{n\mathbf{k}}}{\partial \omega} \right|_{\omega=z/\hbar} (\hbar\omega - z) + \dots$$

Quasiparticle shift and broadening

Replace the Taylor expansion inside the spectral function and rearrange:

$$A(\mathbf{k}, \omega) = \sum_n Z_{n\mathbf{k}} \frac{1}{\pi} \frac{\Gamma_{n\mathbf{k}}}{(\hbar\omega - E_{n\mathbf{k}})^2 + \Gamma_{n\mathbf{k}}^2}$$

Quasiparticle shift and broadening

Replace the Taylor expansion inside the spectral function and rearrange:

$$A(\mathbf{k}, \omega) = \sum_n Z_{n\mathbf{k}} \frac{1}{\pi} \frac{\Gamma_{n\mathbf{k}}}{(\hbar\omega - E_{n\mathbf{k}})^2 + \Gamma_{n\mathbf{k}}^2}$$

$$E_{n\mathbf{k}} = \varepsilon_{n\mathbf{k}} + \text{Re } \Sigma_{n\mathbf{k}}(E_{n\mathbf{k}}/\hbar) \quad \text{quasiparticle energy}$$

Quasiparticle shift and broadening

Replace the Taylor expansion inside the spectral function and rearrange:

$$A(\mathbf{k}, \omega) = \sum_n Z_{n\mathbf{k}} \frac{1}{\pi} \frac{\Gamma_{n\mathbf{k}}}{(\hbar\omega - E_{n\mathbf{k}})^2 + \Gamma_{n\mathbf{k}}^2}$$

$$E_{n\mathbf{k}} = \varepsilon_{n\mathbf{k}} + \text{Re } \Sigma_{n\mathbf{k}}(E_{n\mathbf{k}}/\hbar) \quad \text{quasiparticle energy}$$

$$\Gamma_{n\mathbf{k}} = Z_{n\mathbf{k}} \text{Im } \Sigma_{n\mathbf{k}}(E_{n\mathbf{k}}/\hbar) \quad \text{quasiparticle broadening}$$

Quasiparticle shift and broadening

Replace the Taylor expansion inside the spectral function and rearrange:

$$A(\mathbf{k}, \omega) = \sum_n Z_{n\mathbf{k}} \frac{1}{\pi} \frac{\Gamma_{n\mathbf{k}}}{(\hbar\omega - E_{n\mathbf{k}})^2 + \Gamma_{n\mathbf{k}}^2}$$

$$E_{n\mathbf{k}} = \varepsilon_{n\mathbf{k}} + \text{Re } \Sigma_{n\mathbf{k}}(E_{n\mathbf{k}}/\hbar)$$

quasiparticle energy

$$\Gamma_{n\mathbf{k}} = Z_{n\mathbf{k}} \text{Im } \Sigma_{n\mathbf{k}}(E_{n\mathbf{k}}/\hbar)$$

quasiparticle broadening

$$Z_{n\mathbf{k}} = \frac{1}{1 - \frac{1}{\hbar} \left. \frac{\partial \text{Re } \Sigma_{n\mathbf{k}}(\omega)}{\partial \omega} \right|_{\omega=E_{n\mathbf{k}}/\hbar}}$$

quasiparticle strength

The mass enhancement parameter

Take \mathbf{k} -derivatives of the quasiparticle energy $E_{n\mathbf{k}}$
to find **velocity** and **mass renormalization**[†]

$$V_{n\mathbf{k}} = \frac{v_{n\mathbf{k}}}{1 + \lambda_{n\mathbf{k}}} \quad M_{n\mathbf{k}}^* = (1 + \lambda_{n\mathbf{k}}) m_{n\mathbf{k}}^*$$

[†]These expressions are for the electron gas; more complex expressions are needed in other cases

The mass enhancement parameter

Take \mathbf{k} -derivatives of the quasiparticle energy $E_{n\mathbf{k}}$
to find **velocity** and **mass renormalization**[†]

$$V_{n\mathbf{k}} = \frac{v_{n\mathbf{k}}}{1 + \lambda_{n\mathbf{k}}} \quad M_{n\mathbf{k}}^* = (1 + \lambda_{n\mathbf{k}}) m_{n\mathbf{k}}^*$$

$\lambda_{n\mathbf{k}}$ is the **mass enhancement parameter**

$$\lambda_{n\mathbf{k}} = \frac{1}{Z_{n\mathbf{k}}} - 1 = -\frac{1}{\hbar} \left. \frac{\partial \text{Re}\Sigma_{n\mathbf{k}}(\omega)}{\partial \omega} \right|_{\omega=E_{n\mathbf{k}}/\hbar}$$

[†]These expressions are for the electron gas; more complex expressions are needed in other cases

Electron lifetimes

$$\tau_{n\mathbf{k}} = \frac{\hbar}{2\Gamma_{n\mathbf{k}}} = \frac{\hbar}{2|Z_{n\mathbf{k}} \text{Im} \Sigma_{n\mathbf{k}}(E_{n\mathbf{k}}/\hbar)|}$$

Electron lifetimes

$$\tau_{n\mathbf{k}} = \frac{\hbar}{2\Gamma_{n\mathbf{k}}} = \frac{\hbar}{2|Z_{n\mathbf{k}} \text{Im } \Sigma_{n\mathbf{k}}(E_{n\mathbf{k}}/\hbar)|}$$

Common **approximation**: replace $E_{n\mathbf{k}}$ by $\varepsilon_{n\mathbf{k}}$ and set $Z_{n\mathbf{k}} = 1$

Electron lifetimes

$$\tau_{n\mathbf{k}} = \frac{\hbar}{2\Gamma_{n\mathbf{k}}} = \frac{\hbar}{2|Z_{n\mathbf{k}} \text{Im } \Sigma_{n\mathbf{k}}(E_{n\mathbf{k}}/\hbar)|}$$

Common **approximation**: replace $E_{n\mathbf{k}}$ by $\varepsilon_{n\mathbf{k}}$ and set $Z_{n\mathbf{k}} = 1$

$$\begin{aligned} \frac{1}{\tau_{n\mathbf{k}}} &= \frac{2\pi}{\hbar} \sum_{m\nu} \int_{\Omega_{\text{BZ}}} \frac{d\mathbf{q}}{|g_{nm\nu}(\mathbf{k}, \mathbf{q})|^2} \\ &\quad \times [(1 - f_{m\mathbf{k}+\mathbf{q}} + n_{\mathbf{q}\nu})\delta(\varepsilon_{n\mathbf{k}} - \hbar\omega_{\mathbf{q}\nu} - \varepsilon_{m\mathbf{k}+\mathbf{q}}) \quad \text{phonon emission} \\ &\quad + (f_{m\mathbf{k}+\mathbf{q}} + n_{\mathbf{q}\nu})\delta(\varepsilon_{n\mathbf{k}} + \hbar\omega_{\mathbf{q}\nu} - \varepsilon_{m\mathbf{k}+\mathbf{q}})] \quad \text{phonon absorption} \end{aligned}$$

Identical to Fermi Golden rule formula discussed in Lec. Mon.1

Example: Mass enhancement and lifetimes in MAPbI_3






Figure adapted from Schlipf et al, Phys. Rev. Lett. 121, 086402 (2018)

Take-home messages

- Field theory provides a rigorous and systematic framework for studying electron-phonon physics
- The electron-phonon self-energy yields the mass enhancement and the lifetimes resulting from EPIs
- Many-body formulas are similar but not identical to elementary perturbation theory

References

- FG, Rev. Mod. Phys. 89, 015003 (2017) [\[link\]](#)
- S. Engelsberg and J. R. Schrieffer, Phys. Rev. 131, 993 (1963) [\[Link\]](#)
- A. Eiguren, C. Ambrosch-Draxl, and P. M. Echenique, Phys. Rev. B 79, 245103 (2009) [\[Link\]](#)
- P. Vogl, Phys. Rev. B 13, 694 (1976) [\[Link\]](#)
- E. G. Maksimov, Sov. Phys. JETP 42, 1138 (1976) [\[Link\]](#)
- G. Grimvall, *The electron-phonon interaction in metals*, 1981, (North-Holland, Amsterdam)
- Abrikosov et al, *Methods of quantum field theory in statistical physics*, 1964
- L. Hedin and S. Lundqvist, *Effects of electron-electron and electron-phonon interactions on the one-electron states of solids*, Ed. Seitz, Turnbull, and Ehrenreich, Solid State Physics, Vol. 23 (Academic, 1969)